

Diagonal Ramsey numbers in multipartite graphs related to stars

Chula Jayawardene

Department of Mathematics
University of Colombo, Sri Lanka.

c_jayawardene@yahoo.com

Abstract

Abstract: Let the star on n vertices, namely $K_{1,n-1}$ be denoted by S_n . If every two coloring of the edges of a complete balanced multipartite graph $K_{j \times s}$ there is a copy of S_n in the first color or a copy of S_m in the second color, then we will say $K_{j \times s} \rightarrow (S_n, S_m)$. The size Ramsey multipartite number $m_j(S_n, S_m)$ is the smallest natural number s such that $K_{j \times s} \rightarrow (S_n, S_m)$. In this paper, we obtain the **exact** values of the size Ramsey numbers $m_j(S_n, S_m)$ for $n, m \geq 3$ and $j \geq 3$.

1 Introduction

In this paper we concentrate on simple graphs without loops and multiple edges. Let the complete multipartite graph having j uniform sets of size s be denoted by $K_{j \times s}$ and the complete bipartite graph on $n + m$ vertices be denoted by $K_{n,m}$. Given, three graphs K_N , G and H , we say that $K_N \rightarrow (G, H)$ if K_N is colored by two colors red and blue and it contains a copy of G (in the first color) or a copy of H (in the second color). Using this notation we define the classical Ramsey number $r(n, m)$ as the smallest integer N such that $K_N \rightarrow (K_n, K_m)$. Sadly to say, even in the case of diagonal classical Ramsey numbers $r(n, n)$ almost nothing significant is known beyond the case $n = 5$ (see Radziszowski et al 2014 for a survey).

In the decades that followed there are several interesting variations that have originated from these classical Ramsey number. One obvious variation is the case of size Ramsey numbers build up mainly by Erdős, Faudree, Rousseau and Shelp (see Erdős et al 1978 and Faudree et al 1975). Another variation introduced recently by Buger et al 2004 and Syafrizal et al 2005, is the concept of balanced multipartite Ramsey numbers. This concept is based on exploring the two colorings of multipartite graphs $K_{j \times s}$ instead